

Brad's* parents brought him to see me when he was eleven years old. He was scratched by a cat at age two and developed enlarged lymph nodes in his neck. As a toddler, he began experiencing episodes of inconsolable crying and rocking. By age five, he clearly had problems focusing and was not interested in reading.



Photo by Anna Shvets

His symptoms worsened around age eight. The boy became agitated and easily overwhelmed, with outbursts of anger and rage. His father often had to hold Brad so that he didn't destroy the house.

Brad described dark episodes when he had obsessive, intrusive thoughts, such as hurting himself and others. He had severe anxiety and refused to go to school. He had pseudo-seizures, when he would suddenly flop to the ground but be totally conscious.

Brad had other symptoms, including sharp chest pains with shortness of breath; abdominal pain with nausea, constipation, and flatulence; knee, neck, and spine pain; lightheadedness; sound sensitivity; fatigue; and sugar cravings.

His mood issues were so severe that he underwent two psychiatric hospitalizations. His psychiatrist tried several medications. There was some improvement on Prozac and Trileptal, an anti-epileptic drug that is also prescribed as a mood stabilizer.

We'll get back to Brad. But first, I want to describe one way that microbes can cause psychiatric symptoms.

**Patient's name has been changed to protect his privacy.*

PANS and PANDAS



Dr. Susan Swedo is a researcher in the field of pediatrics and neuropsychiatry. Since 1998, she has been chief of the Pediatrics & Developmental Neuroscience Branch at the U.S. National Institute of Mental Health.

In 1994, Swedo was the lead author on a paper characterizing PANDAS — pediatric autoimmune neuropsychiatric disorders associated with streptococcus. Swedo described children who experienced the sudden onset of obsessive-compulsive disorder, tics, and other mood and behavior problems following a strep infection.¹

Since then, it has become clear that strep is not the only microbe that can trigger psychiatric symptoms. Accordingly, this syndrome is now referred to as PANS — pediatric acute-onset neuropsychiatric syndrome.

Herpes simplex virus, influenza A virus, varicella zoster virus, Epstein-Barr virus, HIV, recurrent sinusitis, mold, and the common cold have been identified as triggers.²⁻⁴ Tick-borne infections, including *Mycoplasma pneumonia*, *Bartonella henselae*, and probably *Borrelia burgdorferi*, the Lyme pathogen, have also been identified as triggers.^{2,4,5} PANDAS, caused by strep, is now considered a subset of PANS.

PANS may be a lot more common than we realize. In 2018, Dr. Nancy Brown and I did a pilot study in which we tested 10 adolescents at a residential treatment center in Colorado.⁶ None of the teens had a known organic illness. All 10 were diagnosed with major depressive disorder and 7 with generalized anxiety disorder. They were too symptomatic to stay at home or go to public school.

Children with PANS often fall behind developmentally and academically. Other common symptoms include involuntary movements, tics, and eating disorders.

[Click to tweet](#)

According to testing, 6 of the 10 kids had antibodies to tick-borne infections—Lyme disease, bartonellosis, and tick-borne relapsing fever. Three had antibodies to *Streptococcus*. What was most surprising was that 9 of the 10 teens had elevated antineuronal antibodies.

Elevation of Antineuronal Antibodies in PANS



Elevation of these antibodies, along with activation of a specific enzyme (the calcium calmodulin-dependent protein kinase II, or CaMKII) is the laboratory hallmark of PANS.⁷

In the 1990s, Dr. Madeleine Cunningham and colleagues performed groundbreaking research to demonstrate the association of these lab abnormalities in kids with this disorder.⁸

It turns out that antibodies that target some microbes attack neuronal tissue and activate

CaMKII, which disrupts dopamine transmission.⁹ The result is “brain on fire”—neuroinflammation that triggers a host of neuropsychiatric symptoms.

Children with PANS display a wide array of mood and behavior issues, especially anxiety, obsessive-compulsive disorder, depression, and irritability/oppositional/anger issues, which can reach epic proportions like Brad’s. Interestingly, many outbursts are followed by remorse.

Children with PANS often fall behind developmentally and academically. Other common symptoms include involuntary movements, tics, and eating disorders. While the case definition of PANS limits the diagnosis to symptoms that have a sudden onset,¹⁰ most of my patients’ symptoms have come on gradually.

In my medical practice (which is limited to patients with tick-borne infections), I see many children and teens. I have been surprised by how many suffer from PANS. While they all have Lyme disease, it appears that Bartonella and Mycoplasma may be the worst offenders in triggering PANS.

What Is the Best Treatment for PANS?

It is important that these kids be treated with antibiotics,¹¹ although careful titration is necessary because Herxheimer reactions can be severe.



Dr. Amiram Katz, a neurologist in Connecticut, has found that low-dose penicillin injections have been very effective.¹² I have had the same experience. Katz recommends long-acting penicillin injections (LA Bicillin), 600,000 units once weekly if the patient weighs less than 100 lbs, and 1.2 million units once weekly if more than 100 lbs. If there is no significant improvement after 10 doses, then there is no reason to continue.

Intravenous gammaglobulin (IVIG) has helped many kids with PANS.¹³ IVIG is composed of human antibodies and has the paradoxical function of stimulating the immune system while decreasing inflammation. It is administered intravenously or subcutaneously, often every three weeks. However, it is quite expensive and insurance coverage is difficult to procure—except for residents of Illinois.

Other interventions are often helpful. Many kids with PANS have mast cell activation syndrome (MCAS), which contributes to inflammation.¹⁴ In these cases, interventions such as antihistamines can be helpful. Cromolyn, which inhibits the release of histamine and other inflammatory mediators from mast cells, is particularly beneficial.

Problem Foods Can Trigger Inflammation

Food sensitivities can also contribute to inflammation,¹⁵ so parents should consider specialized testing or an elimination/challenge diet. Be particularly suspicious of food cravings—the foods people crave are often the worst offenders.¹⁶

In my practice, sugar, yeast, gluten, and dairy products are the most common culprits. Mold can trigger inflammation as well. Even though I live in Colorado, a low-humidity state, I am amazed at how many of my patients suffer from mold issues.

Other interventions are occasionally beneficial. Anti-inflammatory agents, whether ibuprofen or natural products such as curcumin, may help. Some people find that low-dose naltrexone decreases symptoms. Family counseling and support groups are also a good idea. Occasionally, antidepressants—particularly sertraline—can lessen symptoms.

These kids are suffering a great deal. If we do careful detective work and treat the sources of their inflammation as well as the inflammation itself, we can help them get their lives back.

Other modalities that sometimes prove helpful include biofeedback and plasmapheresis, a procedure in which blood is removed and antibodies are filtered out before the blood is replaced.¹²

Back to Brad

Brad suffered from several tick-borne infections: Lyme disease, babesiosis, and bartonellosis. He likely acquired Bartonella from the cat scratch at age two, which then activated a congenital infection with *Borrelia burgdorferi* and Babesia.

He is sensitive to gluten, dairy, and yeast, which I suspected from the pattern of his food cravings. He had a positive [Moleculera Labs](#), which performs the Cunningham Panel, offers a host of information on PANS and PANDAS.

Dr. Daniel Kinderlehrer is an internal medicine physician with a private practice in Denver, Colorado, devoted to treating patients with tick-borne illness. He is the author of the forthcoming book Recovery From Lyme: The Integrative Medicine Guide to the Diagnosis and Treatment of Tick-Borne Illness, which will be released next year.

References:

1. Swedo SE, Leonard HL, Kiessling LS. Speculations on Antineuronal Antibody-Mediated Neuropsychiatric Disorders of Childhood. *Pediatrics*. 1994Feb1;93(2):323-6.
2. Frankovich J, Thienemann M, Rana S, Chang K. Five Youth with Pediatric Acute-Onset Neuropsychiatric Syndrome of Differing Etiologies. *J Child Adolesc Psychopharmacol*. 2015;25(1):31-7. DOI: [10.1089/cap.2014.0056](https://doi.org/10.1089/cap.2014.0056).
3. Tisi G, Marzolini M, Biffi G. Pediatric acute onset neuropsychiatric syndrome associated with Epstein-Barr infection in child with Noonan syndrome. *Europ Psychiatry*. 2017;41(Supplement):S456. DOI: [10.1016/j.eurpsy.2017.01.492](https://doi.org/10.1016/j.eurpsy.2017.01.492).
4. Breitschwerdt EB, Greenberg R, Maggi RG, Mozayeni BR, Lewis A, Bradley JM. Bartonella henselae Bloodstream Infection in a Boy With Pediatric Acute-Onset Neuropsychiatric Syndrome. *J Cent Nerv Syst Dis*. 2019Mar18;11. DOI: [10.1177/1179573519832014](https://doi.org/10.1177/1179573519832014).
5. Rhee H, Cameron D. Lyme disease and pediatric autoimmune neuropsychiatric disorders associated with streptococcal infections (PANDAS): an overview. *Int J Gen Med*. 2012;5:163-74. DOI: [10.2147/ijgm.s24212](https://doi.org/10.2147/ijgm.s24212).
6. Kinderlehrer DA, Brown N. Microbial Induced Autoimmune Inflammation as a Cause of Mental Illness in Adolescents: A Pilot Study. Submitted for publication Feb 25, 2020.
7. Shimasaki C, Frye RE, Trifiletti R, Cooperstock M, Kaplan G, Melamed I, et al. Evaluation of the Cunningham Panel™ in pediatric autoimmune neuropsychiatric disorder associated with streptococcal infection (PANDAS) and pediatric acute-onset neuropsychiatric syndrome (PANS): Changes in antineuronal antibody titers parallel changes in patient symptoms. *J Neuroimmunol*. 2020 Feb15;339:577138. DOI: [10.1016/j.jneuroim.2020.577138](https://doi.org/10.1016/j.jneuroim.2020.577138).

[10.1016/j.jneuroim.2019.577138](https://doi.org/10.1016/j.jneuroim.2019.577138).

8. Cunningham MW, Cox CJ. Autoimmunity against dopamine receptors in neuropsychiatric and movement disorders: a review of Sydenham chorea and beyond. *Acta Physiol*. 2016Jan;216(1):90–100. DOI: [10.1111/apha.12614](https://doi.org/10.1111/apha.12614)
9. Cox CJ, Zuccolo AJ, Edwards EV, Mascaro-Blanco A, Alvarez K, Stoner J, et al. Antineuronal Antibodies in a Heterogeneous Group of Youth and Young Adults with Tics and Obsessive-Compulsive Disorder. *J Child Adolesc Psychopharmacol*. 2015Feb;25(1):76–85. DOI: [10.1089/cap.2014.0048](https://doi.org/10.1089/cap.2014.0048).
10. Chang K, Frankovich J, Cooperstock M, Cunningham MW, Latimer ME, Murphy TK, et al. Clinical Evaluation of Youth with Pediatric Acute-Onset Neuropsychiatric Syndrome (PANS): Recommendations from the 2013 PANS Consensus Conference. *J Child Adolesc Psychopharmacol*. 2015;25(1):3–13. DOI: [10.1089/cap.2014.0084](https://doi.org/10.1089/cap.2014.0084).
11. Calaprice D et al. Treatment of Pediatric Acute-Onset Neuropsychiatric Disorder in a Large Survey Population. *J Child Adolesc Psychopharmacol*. 2018;28(2):92–103. DOI: [10.1089/cap.2017.0101](https://doi.org/10.1089/cap.2017.0101)
12. Katz A. (2018, November). NADAL—Neuropsychiatric Autoimmune Disorder Associated With Lyme Disease –A PANDAS/PANS Equivalent: Diagnosis and Treatment. Presented at the 19th annual conference of *The International Lyme and Associated Diseases conference*, Chicago, IL.
13. Kovacevic M, Grant P, Swedo SE. Use of intravenous immunoglobulin in the treatment of twelve youths with pediatric autoimmune neuropsychiatric disorders associated with streptococcal infections. *J Child Adolesc Psychopharmacol*. 2015;25(1):65–69. DOI: [10.1089/cap.2014.0067](https://doi.org/10.1089/cap.2014.0067)
14. Walker ME, Hatfield JK, Brown MA. New insights into the role of mast cells in autoimmunity: evidence for a common mechanism of action? *Biochim Biophys Acta*. 2012;1822(1):57–65. DOI: [10.1016/j.bbadis.2011.02.009](https://doi.org/10.1016/j.bbadis.2011.02.009)
15. Rosa JS et al. Allergic Diseases and Immune-Mediated Food Disorders in Pediatric Acute-Onset Neuropsychiatric Syndrome. *Pediatr Allergy Immunol Pulmonol*. 2018 Sep 1;31(3):158–165. DOI: [10.1089/ped.2018.0888](https://doi.org/10.1089/ped.2018.0888)
16. Randolph TG, Moss RW. *An Alternative Approach to Allergies*. Harper & Row, New York, ©1979.

Editor’s note: Any medical information included is based on a personal experience. For questions or concerns regarding health, please consult a doctor or medical professional.